





HOW TO USE AUVI-Q® (EPINEPHRINE INJECTION, USP), KALEO

1. Remove Auvi-Q from the outer case.
2. Pull off red safety guard.
3. Place black end of Auvi-Q against the middle of the outer thigh.
4. Press firmly, and hold in place for 5 seconds.
5. Call 911 and get emergency medical help right away.




HOW TO USE EPIPEN® AND EPIPEN JR® (EPINEPHRINE) AUTO-INJECTOR, MYLAN

1. Remove the EpiPen® or EpiPen Jr® Auto-Injector from the clear carrier tube.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward.
3. With your other hand, remove the blue safety release by pulling straight up.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'.
5. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
6. Remove and massage the injection area for 10 seconds.
7. Call 911 and get emergency medical help right away.




HOW TO USE EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF EPIPEN®), USP AUTO-INJECTOR, MYLAN

1. Remove the epinephrine auto-injector from the clear carrier tube.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward.
3. With your other hand, remove the blue safety release by pulling straight up.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'.
5. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
6. Remove and massage the injection area for 10 seconds.
7. Call 911 and get emergency medical help right away.



HOW TO USE IMPAX EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF ADRENALCLICK®), USP AUTO-INJECTOR, IMPAX LABORATORIES

1. Remove epinephrine auto-injector from its protective carrying case.
2. Pull off both blue end caps: you will now see a red tip.
3. Grasp the auto-injector in your fist with the red tip pointing downward.
4. Put the red tip against the middle of the outer thigh at a 90-degree angle, perpendicular to the thigh.
5. Press down hard and hold firmly against the thigh for approximately 10 seconds.
6. Remove and massage the area for 10 seconds.
7. Call 911 and get emergency medical help right away.



ADMINISTRATION AND SAFETY INFORMATION FOR ALL AUTO-INJECTORS:

1. Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than mid-outer thigh. In case of accidental injection, go immediately to the nearest emergency room.
2. If administering to a young child, hold their leg firmly in place before and during injection to prevent injuries.
3. Epinephrine can be injected through clothing if needed.
4. Call 911 immediately after injection.

OTHER DIRECTIONS/INFORMATION (may self-carry epinephrine, may self-administer epinephrine, etc.):

Treat the person before calling emergency contacts. The first signs of a reaction can be mild, but symptoms can worsen quickly.

EMERGENCY CONTACTS — CALL 911

RESCUE SQUAD: _____
DOCTOR: _____ PHONE: _____
PARENT/GUARDIAN: _____ PHONE: _____

OTHER EMERGENCY CONTACTS

NAME/RELATIONSHIP: _____
PHONE: _____
NAME/RELATIONSHIP: _____
PHONE: _____

PLACE
PICTURE
HERE

Name: _____ D.O.B.: _____

Allergy to: _____

Weight: _____ lbs. Asthma: Yes (higher risk for a severe reaction) No

NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE EPINEPHRINE.

Extremely reactive to the following allergens: _____

THEREFORE:

If checked, give epinephrine immediately if the allergen was **LIKELY** eaten, for **ANY** symptoms.

If checked, give epinephrine immediately if the allergen was **DEFINITELY** eaten, even if no symptoms are apparent.

FOR ANY OF THE FOLLOWING:
SEVERE SYMPTOMS



LUNG

Shortness of breath, wheezing, repetitive cough



HEART

Pale or bluish skin, faintness, weak pulse, dizziness



THROAT

Tight or hoarse throat, trouble breathing or swallowing



MOUTH

Significant swelling of the tongue or lips



SKIN

Many hives over body, widespread redness



GUT

Repetitive vomiting, severe diarrhea



OTHER

Feeling something bad is about to happen, anxiety, confusion

OR A
COMBINATION
of symptoms
from different
body areas.

1. **INJECT EPINEPHRINE IMMEDIATELY.**
2. **Call 911.** Tell emergency dispatcher the person is having anaphylaxis and may need epinephrine when emergency responders arrive.
- Consider giving additional medications following epinephrine:
 - » Antihistamine
 - » Inhaler (bronchodilator) if wheezing
- Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
- If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
- Alert emergency contacts.
- Transport patient to ER, even if symptoms resolve. Patient should remain in ER for at least 4 hours because symptoms may return.

MILD SYMPTOMS



NOSE

Itchy or runny nose, sneezing



MOUTH

Itchy mouth



SKIN

A few hives, mild itch



GUT

Mild nausea or discomfort

FOR MILD SYMPTOMS FROM MORE THAN ONE SYSTEM AREA, GIVE EPINEPHRINE.

FOR MILD SYMPTOMS FROM A SINGLE SYSTEM AREA, FOLLOW THE DIRECTIONS BELOW:

1. Antihistamines may be given, if ordered by a healthcare provider.
2. Stay with the person; alert emergency contacts.
3. Watch closely for changes. If symptoms worsen, give epinephrine.

MEDICATIONS/DOSES

Epinephrine Brand or Generic: _____

Epinephrine Dose: 0.15 mg IM 0.3 mg IM

Antihistamine Brand or Generic: _____

Antihistamine Dose: _____

Other (e.g., inhaler-bronchodilator if wheezing): _____

Authorization for the Administration of Medication by School, Child Care, and Youth Camp Personnel

In Connecticut schools, licensed Child Day Care Centers and Group Day Care Homes, licensed Family Day Care Homes, and licensed Youth Camps administering medications to children shall comply with all requirements regarding the Administration of Medications described in the State Statutes and Regulations. Parents/guardians requesting medication administration to their child shall provide the program with appropriate written authorization(s) and the medication before any medications are administered. Medications must be in the original container and labeled with child's name, name of medication, directions for medication's administration, and date of the prescription.

Authorized Prescriber's Order (Physician, Dentist, Optometrist, Physician Assistant, Advanced Practice Registered Nurse or Podiatrist):

Name of Child/Student _____ Date of Birth ____/____/____ Today's Date ____/____/____

Address of Child/Student _____ Town _____

Medication Name/Generic Name of Drug _____ Controlled Drug? YES NO

Condition for which drug is being administered: _____

Specific Instructions for Medication Administration _____

Dosage _____ Method/Route _____

Time of Administration _____ If PRN, frequency _____

Medication shall be administered: Start Date: ____/____/____ End Date: ____/____/____

Relevant Side Effects of Medication _____ None Expected

Explain any allergies, reaction to/negative interaction with food or drugs _____

Plan of Management for Side Effects _____

Prescriber's Name/Title _____ Phone Number (____) _____

Prescriber's Address _____ Town _____

Prescriber's Signature _____ Date ____/____/____

School Nurse Signature (if applicable) _____

Parent/Guardian Authorization:

I request that medication be administered to my child/student as described and directed above

I hereby request that the above ordered medication be administered by school, child care and youth camp personnel and I give permission for the exchange of information between the prescriber and the school nurse, child care nurse or camp nurse necessary to ensure the safe administration of this medication. I understand that I must supply the school with no more than a three (3) month supply of medication (school only.)

I have administered at least one dose of the medication with the exception of emergency medications to my child/student without adverse effects. (For child care only)

Parent/Guardian Signature _____ Relationship _____ Date ____/____/____

Parent /Guardian's Address _____ Town _____ State _____

Home Phone # (____) _____ - _____ Work Phone # (____) _____ - _____ Cell Phone # (____) _____ - _____

SELF ADMINISTRATION OF MEDICATION AUTHORIZATION/APPROVAL

Self-administration of medication may be authorized by the prescriber and parent/guardian and must be approved by the school nurse (if applicable) in accordance with board policy. In a school, inhalers for asthma and cartridge injectors for medically-diagnosed allergies, students may self-administer medication with only the written authorization of an authorized prescriber and written authorization from a student's parent or guardian or eligible student.

Prescriber's authorization for self-administration: YES NO _____
Signature Date

Parent/Guardian authorization for self-administration: YES NO _____
Signature Date

School nurse, if applicable, approval for self-administration: YES NO _____
Signature Date

Today's Date _____ Printed Name of Individual Receiving Written Authorization and Medication _____

Title/Position _____ Signature (in ink or electronic) _____

Note: This form is in compliance with Section 10-212a, Section 19a-79-9a, 19a-87b-17 and 19-13-B27a(v.)

